

Comparison of Oral Health-related Quality of Life in Treated and Untreated Class II Adolescent Patients

Fahimeh Farzanegan¹, Farnaz Zia², Lohrasb Dehghani^{2*}, Milad Zarei²

¹Professor, Dental Research Center, Mashhad University of Medical Sciences, Mashhad, Iran ²Resident, Department of Orthodontics, Mashhad University of Medical Sciences, Mashhad, Iran

*Corresponding author: Lohrasb Dehghani

Address: Faculty of Dentistry, Mashhad University of Medical Sciences, Mashhad, Iran.

Email: lohrasbbio@gmail.com

Received: 2023 May 27; Revised: 2023 October 10; Accepted: 2023 October 31

Abstract

Aim: This study aimed to investigate the effect of growth modification treatments on oral health-related quality of life in adolescents aged 11 to 14 years with Class II malocclusion.

Methods: Eighty people participated in this case-control study. The case group (n=43) included adolescents with a history of Class II malocclusion treated with growth modification. The control group consisted of 37 adolescents with Class II malocclusion who were not treated. Both groups were given a translation of the Child Perception Questionnaire (CPQ), measuring the quality of life in four domains: oral symptoms, functional limitations, psychological health, and social wellbeing. Data were statistically analyzed by SPSS software and a significance level of 0.05 was considered.

Results: The oral health-related quality of life of the case group was better than the control group in all areas although this difference was not statistically significant (P=0.204).

Conclusion: The oral health-related quality of life between two groups was not different; however, the difference in mental health components between the two groups was clinically negligible.

Keywords: Quality of Life, Malocclusion, Angle Class II, Adolescent, Oral Health

Background

According to WHO, the definition of health not only refers to the lack of disease but also to possess adequate health physically, psychologically, and socially. Therefore, in a study, all aspects affecting health should be considered (1).

Malocclusion is one of the most prevalent oral and dental problems, and in terms of WHO ranks third in common oral diseases after caries and periodontal diseases (1). Clear malocclusion, increased overjet, and crowded incisors are related to low self-esteem in adolescents (2). Optimal appearance is also one of the factors in improving socio-psychological status (3). The second most important factor in the appearance of the face after the eyes is the mouth and teeth and the overall appearance of the teeth and face affects the mental image of the person (body image) and can help one's satisfaction, and the higher satisfaction of the person with their overall facial structure, would be the higher the confidence (4).

The results of a study have shown that malocclusion severity could negatively impact the quality of life. Considering the psychological effects of malocclusion on quality of life, clinicians should place more emphasis on the treatment of patients with malocclusions (2, 3). Developmental modification treatments should be performed before or during the growth spurt during adolescence because, after this period, skeletal growth has nearly ceased. In class II malocclusion, depending on which jaw has growth defects or overgrowth, treatments are usually divided into two types of functional devices (stimulating mandible growth) and a variety of headgears (preventing maxillary overgrowth) (4). In cases where dental defects or growth spurts have occurred and growth shaving treatments can no longer be used, fixed treatment is chosen (4).

Copyright © 2023, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

However, to choose the appropriate method, all of the patient's conditions should be considered and the final treatment plan should be chosen according to the clinical conditions and psychosocial conditions of the patient (4,1). The definition of the oral health-related quality of life (OHRQOL) concept, according to the U.S. Public Health Service, is a "structurally multi-dimensional quality of life" and indicates people's ease when eating and sleeping, how much they participate in social interactions, self-esteem, and contentment with their oral health (5).

In class II malocclusion, as mentioned above, there are two main therapeutic pathways. Considerable research has been conducted on the fixed path (6, 7) and a positive correlation exists between fixed treatments and OHRQOL (8). However, oral quality of life in removable aligners is less evident and there is no accurate comparison between these two treatments. According to studies, during fixed orthodontic treatment, especially in the early months, there is a decrease in quality of life, but over time this index improves (9). Due to the lack of investigation of this issue in different studies, this study aimed to investigate the effects of growth modification treatments on oral health-related quality of life in adolescents aged 11 to 14 years with class II malocclusion.

Methods

Eighty patients participated in this study. The case group was composed of 43 patients between 11 and 14 years old who were treated by class II growth change treatments including functional and headgear devices in the orthodontics department of the Mashhad Dental School and had 1-3 mm overjet and half-face improvement. (The overjet of the case group was greater than or equal to 5 mm before treatment.)

Parallel with this group, the control group consisted of 37 untreated patients with grade II (overjet greater than or equal to 5 mm and convex protrusion), with the same age range of 11 to 14 years and with the same sex dispersion. These subjects were selected from students of Nasim Vahid Girls' School located on Majd Boulevard and patients referred to the orthodontic department of the faculty for which treatment had not yet begun. Both groups were given a 14 CPQ11 questionnaire, wherein the orthodontics department authenticated its validity and reliability (Cronbach's alpha coefficient was 0.894 and ICC was 0.91), and according to the obtained scores, the quality of liferelated to the oral health of these adolescents was measured and compared.

To evaluate the distribution of quantitative variables, the Shapiro–Wilk test was used. To investigate the influential variables such as age, overjet, and sex, the Mann-Whitney and chi-square tests were used.

The scores of the domains of the two groups were analyzed and compared using the independent t-tests, and Mann-Whitney odor tests were used. The statistical significance level was set to 5%.

Results

In this study, the quality of life in the two groups of students was assessed using the CPQ11-14 questionnaire in 2021 in general and partially (in different domains). The total participants were 80 (case group: 43 subjects, control group: 37 subjects), of whom 66.3% were female and 33.7% were male. The mean and standard deviation of age was 11.8±1.7 years and their age range was 11 to 14 years.

Next, we described the scores of all domains and the overall score of the questionnaire and compared the scores obtained in each area and the overall score of the questionnaire in terms of the group. The overall status of the score of each domain and the total score are as follows: The highest and lowest mean domains were related to "oral problems" and "functional limitations", respectively. The lowest and highest total scores were 0 and 77, respectively. The mean and median of the total scores were 25.79 and 23, respectively.

Then the normality of the distribution of age, overjet, and each domain and total score in the studied groups were evaluated using the Shapiro-Wilk test. The results indicated that only the scores of oral problems in both groups had normal distribution, so this content was considered in the data analysis (Table 1).

The variables affecting age, sex, and overjet in the two groups were also compared. In Table 2, the comparison of age and overjet between groups is shown. In Table 3, the frequency distribution of sex in the studied groups is shown. No significant difference existed between the groups regarding confounding variables. In this section, we compared the total score and scores of each domain between the groups.

Table 1. Shapiro-Wilk test results for normalization of data distribution of quantitative variables studied

Dehghani L et al.

Variable		Statistics	Degrees of Freedom	P value	Distribution
Ago.	Case	0.942	43	0.032	Abnormal
Age	Control	0.918	37	0.010	Abnormal
Overiet	Case	0.897	43	0.001	Abnormal
Overjet	Control	0.833	37	0.000	Abnormal
Oral Brobloms	Case	0.973	43	0.392	Normal
Oral Problems	Control	0.942	37	0.054	Normal
Functional Limitations	Case	0.882	43	<0.001	Abnormal
Functional Limitations	Control	0.870	37	<0.001	Abnormal
Emotional and Montal Health	Case	0.838	43	<0.001	Abnormal
	Control	0.907	37	0.005	Abnormal
Social Walfara	Case	0.703	43	< 0.001	Abnormal
Social Wellare	Control	0.807	37	<0.001	Abnormal
Total Score	Case	0.902	43	0.001	Abnormal
	Control	0.905	37	0.004	Abnormal

The score of each domain and the total score of the questionnaire between the groups were compared. In Table 4, the score of each domain and the overall score of the questionnaire between the groups are given. In Table 5, the relationship between age and domains and overall score is given. It can be observed that in the case group, only the domain of "functional constraints" had a direct relationship (P=0.016). Other domains and overall scores indicated no significant association with age.

Furthermore, other domains and total scores were not significantly correlated with overjet. In the control group, none of the domains and total score had a significant relationship with any of the variables of age and overjet.

Table 2. Comparison of age and overjet between groups							
Variable	Group	Number	Standard Deviation±Average	Middle	Most Least		Mann-Whitney U result
A.c.o.	Case	43	12.4±1.6	12.0 (3.0)	14	11	Z=2.90
Age	Control	37	11.2±1.7	11.0 (3.0)	14	11	P=0.0 [≤] •
Overiet	Case	43	7.4±2.3	7.0 (4.0)	14	5	Z=0.75
Overjet	Control	37	7.1±2.4	6.0 (4.0)	14	5	P=0.452

Table 3. Frequency distribution of sex in the studied groups

			Se	Total		
			Female Male		iotai	
	Casa	Number	31	12	43	
Crown	Case	%	72.1%	72.9%	100.0%	
Group	Control	Number	22	15	37	
		%	59.5%	40.5%	100.0%	
Total		Number	53	27	80	
TOLAI		%	66.3%	33.8%	100.0%	
Chi-square result		X2=1.42 p=0.233				

 Table 4. Comparison of scores in each domain and total score between groups

Area	Sex	Number	Standard Deviation±Average	Middle	Most Least	Mann-Whitney Result
Oral Problems	Case	43	7.2±4.2	7.0 (6.0)	16 0	T=1.25
Oral FTODIETTIS	Control	37	8.5±5.2	7.0 (8.0)	22 0	P=0.216
Functional	Case	43	4.9±4.6	4.0 (6.0)	18 0	Z=0.07
Limitation	Control	37	5.1±5.2	4.0 (7.5)	19 0	P=0.942
Emotional and	Case	43	5.2±5.3	4.0 (7.0)	20 0	Z=1.65
Mental Health	Control	37	7.3±6.2	6.0 (8.0)	26 0	P=0.100
Social Welfare	Case	43	6.1±7.8	3.0 (5.0)	34 0	Z=0.90
	Control	37	7.6±8.2	4.0 (9.0)	30 0	P=0.369
Total Score	Case	43	23.4±16.0	21.0 (21.0)	77 0	Z=1.27
	Control	37	28.5±19.1	23.0 (22.0)	74 0	P=0.204

T: Independent Samples t-Test

Table 5. The relationship between age and overjet with domains and total score								
Group	Variable		Oral Problems	Functional Limitation	Emotional and Mental Health	Social Welfare	Total Score	
Case (number=4) Control (number=3)	Age	Spearman's rank correlation coefficient	0.151	0.367	0.099	0.065	0.21	
	Overiet	P value Spearman's rank correlation	0.335 0.115	0.016 0.173	0.527	0.677 0.169	0.176 0.259	
	Overjet	coefficient P value	0.462	0.268	0.576	0.279	0.094	
	Age	correlation coefficient	-0.212	-0.155	0.017	0.15	0.018	
		P value Spearman's rank	0.207	0.36	0.921	0.375	0.918	
	Overjet	correlation coefficient P value	-0.234 0.163	0.145	-0.027 0.875	-0.092	-0.105 0.537	

Discussion

People with malocclusions and dental deformities may experience physical, social, and psychological difficulties. Malocclusion treatment can be used efficiently to improve function, facial appearance, and dental aesthetics. Generally, patients report their motivations for seeking treatment are the functional and aesthetic factors or self-confidence and self-esteem; however, the number of patients reporting these motives varies widely between studies.

This study aimed to evaluate the effect of orthodontic growth modification treatments on the quality of life components of adolescents. In this regard, two groups between the ages of 11 and 14 years answered the CPQ questionnaire. The first group (case) consisted of the subjects who had a history of class II malocclusion (overjet more than 5 mm) and were at that time undergoing treatment. The second group (control) consisted of subjects with class II malocclusion and who have not been treated.

By comparing the results, we found out that in the four domains of the questionnaire, which included oral symptoms, functional limitations, emotional and mental health, and social welfare, the control group had higher scores in all components and therefore better quality of life compared to the case group. Although the difference indicated no statistical significance (P=0.204), the mean difference between both groups in the field of mental health is considerable and clinically inconspicuous. The study's outcome was consistent with those of Palomares et al., who identified improved OHRQoL for adults who had finished orthodontic treatment and were in the retention phase versus their non-treated counterparts (10). Chinese orthodontic patients were found to have a better OHQoL after they finished treatment as compared with preorthodontic treatment and during the treatment by Chen et al (11).

A noteworthy point is that the effect of class II malocclusion on quality of life may be less than other types. For example, studies were conducted by Mostafa Quchani (2011) and Aydin Faceger (2014) on adolescent boys and girls ages 14 to 17. The crowding of teeth, especially in the anterior, which has a great impact on esthetic, has more negative effects on adolescents' quality of life (12, 13). However, it may not have a significant effect on a person's body image against class II malocclusion. Usually, people see their image in the mirror from the front, hence, the convex profile does not receive much attention. Therefore, the level of quality of life before treatment and in different types of malocclusions can have a wide spectrum, and an important factor is the effects of malocclusion on the beauty of the patient, it is expected that in adolescents, static is one of the most important and influential factors. However, many studies show that fixed orthodontic treatments regardless of malocclusion have increased quality of life (14).

In a study by Farzanegan et al., decreased quality of life at the beginning of fixed orthodontic treatment was reported, however, it gradually increased and at the end of treatment, significant improvement was observed compared to pretreatment (6). This result falls in the same direction as the Zhang study done in 2008 (15).

Conflicting results include the Taylor Study in 2009 (16). In this study, there was no significant

difference in the quality-of-life components after treatment compared to pre-treatment. Also, in the WC study. Shaw did not find enough evidence of the long-term effect of the need for orthodontic treatment on the mental health of individuals (17). However, in a systematic review article published in 2015 with a review of high-quality articles, the undesirable effect of malocclusion on the quality of life index, especially in psychological and social dimensions, was concluded (18).

Conclusion

This study aimed to show that malformations in the jaws and teeth can harm adolescents' quality of life. Hence, the effects of one of the oral malformations on the quality of life of adolescents 11-14 years old was investigated. The patients with class II malocclusion in the four components of oral symptoms, oral functioning limitation, mental health, and social welfare had a lower quality of life score in the CPQ questionnaire than those treated. Although the difference in the quality of life components had not reached a statistically significant level, the difference between the two groups in the mental health component was clinically significant. The excessively normal front of the maxilla, which causes a convex half-face, can affect adolescents' emotional and mental health. Therefore, paying attention to malocclusion and its appropriate treatment effectively can improve the quality of life in children and adolescents and many studies confirm this. Also, parents should pay special attention in this regard, particularly in case of limitation of function in the oral area (such as difficulties in chewing or speaking, etc.) or cosmetic defects, and their effect on the psychosocial health of children and adolescents. Therefore, it is recommended to go to a general dentist or orthodontist to receive necessary guidance.

References

- Ferrando-Magraner E, Garcia-Sanz V, Bellot-Arcis C, Montiel-Company JM, Almerich-Silla JM, Paredes-Gallardo V. Oral health-related quality of life of adolescents after orthodontic treatment. A systematic review. J Clin Exp Dent. 2019;11(2):194-202. doi: 10.4317/jced.55527. PMID: 30805125.
- Trulsson U, Strandmark M, Mohlin B, Berggren U. A qualitative study of teenagers' decisions to undergo orthodontic treatment with fixed appliance. J Orthod. 2002;29(3):197-204. doi: 10.1093/ortho/29.3.197. PMID: 12218197.
- Birkeland K, Bøe OE, Wisth PJ. Relationship between occlusion and satisfaction with a dental appearance in orthodontically treated and untreated groups. A longitudinal study. Europ J

Orthod. 2000;22(5):509-18. doi: 10.1093/ejo/22.5. 509. PMID: 11105407.

- 4. Amnaie A, Nasiri M, Adib M, Shirvani A. Relationship between self-concept and satisfaction with tooth alignment and demand for orthodontic treatment and professionally determined orthodontic needs. J Isfahan Dent Sch. 2012;. 656-61.
- General Usphsoots, Dental NIO, Research C. Oral health in America: a report of the Surgeon General: US Public Health Service, Department of Health and Human Services; 2000.
- Farzanegan F, Heravi F, Ramezani M. Evaluation of health-related quality of life changes after initial orthodontic treatment. Oral Health Prev Dent. 2015;13(2):143-7. doi: 10.3290/j.ohpd.a33087. PMID: 25431801.
- Ferrando-Magraner E, García-Sanz V, Bellot-Arcís C, Montiel-Company JM, Almerich-Silla JM, Paredes-Gallardo V. Oral health-related quality of life of adolescents after orthodontic treatment. A systematic review. J Clin Exp Dent. 2019;11(2):194-202. doi: 10.4317/jced.55527. PMID: 30805125.
- De Oliveira CM, Sheiham A. Orthodontic treatment, and its impact on oral health-related quality of life in Brazilian adolescents. J Orthod. 2004;31(1):20-7. doi: 10.1179/146531204225011364. PMID: 15071148.
- Jokovic A, Locker D, Stephens M, Kenny D, Tompson B, Guyatt G. Validity and reliability of a questionnaire for measuring child oral-health-related quality of life. J Dent Res. 2002;81(7):459-63. doi: 10.1177/154405910208 100705. PMID: 12161456.
- Palomares NB, Celeste RK, de Oliveira BH, Miguel JA. How does orthodontic treatment affect young adults' oral health-related quality of life?. Am J Orthod Dentofacial Orthop. 2012;141(6):751-8. doi: 10.1016/j. ajodo.2012.01.015. PMID: 22640677.
- Locker D, Jokovic A, Tompson B, Prakash P. Is the Child Perceptions Questionnaire for 11–14-year-olds sensitive to clinical and self-perceived variations in orthodontic status?. Community Dent Oral Epidemiol. 2007;35(3):179-85. doi: 10.1111/j.1600-0528.2006. 00324.x. PMID: 17518964.
- Heravi F, Farzanegan F, Tabatabaee M, Sadeghi M. Do malocclusions affect the oral health-related quality of life?. Oral Health Prev Dent. 2011;9(3):229-33. PMID: 22068178.
- Farzanegan F, Heravi F, Sooratgar A, Dastmalchi P. Evaluation of relationship between oral health-related quality of life and occlusion traits among female adolescents. Dental Res J. 2014;11(6):684-8. PMID: 25540664.
- Chen M, Feng ZC, Liu X, Li ZM, Cai B, Wang DW. Impact of malocclusion on oral health–related quality of life in young adults. Angle Orthod. 2015;85(6):986–91. doi: 10.2319/101714-743.1. PMID: 25531421.
- Zhang M, McGrath C, Hägg U. Changes in oral healthrelated quality of life during fixed orthodontic appliance therapy. Am J Orthod Dentofacial Orthop. 2008;133(1):25-9. doi: 10.1016/j.ajodo.2007.01.024. PMID: 18174067.
- Taylor KR, Kiyak A, Huang GJ, Greenlee GM, Jolley CJ, King GJ. Effects of malocclusion and its treatment on the

quality of life of adolescents. Am J Orthod Dentofacial Orthop. 2009;136(3):382–92. doi: 10.1016/j.ajodo. 2008.04.022. PMID: 19732673.

 Shaw WC, Richmond S, Kenealy PM, Kingdon A, Worthington H. A 20-year cohort study of health gain from orthodontic treatment: psychological outcome. Am J Orthod Dentofacial Orthop. 2007;132(2):146-57. doi: 10.1016/j.ajodo.2007.04.009. PMID: 17693363.

 Dimberg L, Arnrup K, Bondemark L. The impact of malocclusion on the quality of life among children and adolescents: a systematic review of quantitative studies. Eur J Orthod. 2015;37(3):238-47. doi: 10.1093/ejo/cju046. PMID: 25214504.